arXiv:math/9807126v1 [math.RA] 22 Jul 1998

22 July 1998

THE OCTONIONIC EIGENVALUE PROBLEM

Tevian Dray
Department of Mathematics, Oregon State University, Corvallis, OR 97331, USA
tevian@math.orst.edu

Corinne A. Manogue
Department of Physics, Oregon State University, Corvallis, OR 97331, USA
corinne@physics.orst.edu

ABSTRACT

We discuss the eigenvalue problem for 2×2 and 3×3 octonionic Hermitian matrices. In both cases, we give the general solution for real eigenvalues, and we show there are also solutions with non-real eigenvalues.

1. INTRODUCTION

Finding the eigenvalues and eigenvectors of a given matrix is one of the basic techniques in linear algebra, with countless applications. The simplest case is that of (complex) Hermitian matrices, generalizing the familiar case of (real) symmetric matrices. This simple case is nevertheless very important, for instance in quantum mechanics, where the fact that such matrices have real eigenvalues allows them to represent physically observable quantities.

The eigenvalue problem is usually formulated over a field, typically either the real numbers $\mathbb{R}$ or the complex numbers $\mathbb{C}$. We consider here the generalization to the other normed division algebras, namely the quaternions $\mathbb{H}$ and the octonions $\mathbb{O}$. We find that most of the basic properties are retained, provided they are reinterpreted to take into account the lack of commutativity of $\mathbb{H}$ and $\mathbb{O}$, and the lack of associativity of $\mathbb{O}$.

Our main result is the solution of the real eigenvalue problem for 3×3 octonionic Hermitian matrices, also known as Jordan matrices. It is straightforward to show [1] that such matrices admit 24 real eigenvalues, corresponding to eigenvectors which are independent over $\mathbb{R}$. We show that these eigenvalues do not satisfy the characteristic equation even though the matrix itself does. Instead they generically come in 6 sets of multiplicity 4 rather than the expected 3 sets of multiplicity 8 [1]. We further show how to generalize the notion of orthogonality to the nonassociative case, recovering the standard decomposition of a Hermitian matrix in terms of its eigenvalues and eigenvectors.

We begin in Section 2 with a review of the standard eigenvalue problem for real and complex Hermitian matrices, and then consider the quaternionic eigenvalue problem in Section 3. A brief discussion of the properties of octonions and octonionic matrices appears in

[1] After this work was completed, we discovered the existence of earlier work (in Russian) by Ogievetskiĭ [2] which also claims 6 real eigenvalues for such matrices.

Section 4, after which the octonionic eigenvalue problem is considered for 2×2 and 3×3 octonionic Hermitian matrices in Section 5 and Section 6, respectively. Finally, we discuss our results in Section 7.

2. THE STANDARD EIGENVALUE PROBLEM

The eigenvalue problem as usually stated is to find solutions λ, v to the equation

$$Av = \lambda v \tag{1}$$

for a given square matrix A. The basic properties of the eigenvalue problem for $n \times n$ complex Hermitian matrices are well-understood. [2]

Lemma $1_{\mathbb{C}}$: *An $n \times n$ complex Hermitian matrix A has n real eigenvalues (counting multiplicity).*

Proof: We give here only the proof that the eigenvalues are real. Let A, v, λ satisfy (1), with $A^\dagger = A$. Then

$$\overline{\lambda} v^\dagger v = (Av)^\dagger v = v^\dagger A v = \lambda v^\dagger v \tag{2}$$

so that if $v \neq 0$ we have $v^\dagger v \neq 0$, which forces $\overline{\lambda} = \lambda$. ■

Lemma $2_{\mathbb{C}}$: *Eigenvectors of an $n \times n$ complex Hermitian matrix A corresponding to different eigenvalues are orthogonal.*

Proof: For $m = 1, 2$, let v_m be an eigenvector of $A = A^\dagger$ with eigenvalue λ_m. By the previous lemma, $\lambda_m \in \mathbb{R}$. Then

$$\lambda_1 v_1^\dagger v_2 = (Av_1)^\dagger v_2 = v_1^\dagger A v_2 = \lambda_2 v_1^\dagger v_2 \tag{3}$$

Then either $\lambda_1 = \lambda_2$ or $v_1^\dagger v_2 = 0$. ■

Lemma $3_{\mathbb{C}}$: *For any $n \times n$ complex Hermitian matrix A, there exists an orthonormal basis of $\mathbb{C}^n$ consisting of eigenvectors of A.*

Proof: If all eigenvalues have multiplicity one, the result follows from the previous lemma. The Gram-Schmidt orthogonalization process can be used on any eigenspace corresponding to an eigenvalue with multiplicity greater than one. ■

These lemmas are equivalent to the standard result that a complex Hermitian matrix can always be diagonalized by a unitary transformation. It is important for what follows to realize that the form of the proofs given above relies on both the commutativity and the associativity of $\mathbb{C}$.

Combining the above results, it is easy to see that any (complex) Hermitian matrix A admits a decomposition in terms of an orthonormal basis of eigenvectors.

---

[2] We could just as well have started with the case of real symmetric matrices.

Theorem $1_{\mathbb{C}}$: *Let A be an $n \times n$ complex Hermitian matrix. Then A can be expanded as*

$$A = \sum_{m=1}^{n} \lambda_m v_m v_m^\dagger \tag{4}$$

where $\{v_m;\ m = 1, ..., n\}$ is an orthonormal basis of eigenvectors corresponding to eigenvalues λ_m.

Proof: By the previous lemma, there exists an orthonormal basis $\{v_m\}$ of eigenvectors. It then suffices to check that

$$\sum_{m=1}^{n} \lambda_m v_m v_m^\dagger v_k = \lambda_k v_k \tag{5}$$

But this follows by direct computation using orthonormality. ■

Furthermore, the set of eigenvalues $\{\lambda_m\}$ is unique, and the (unit) eigenvectors are unique up to unitary transformations in the separate eigenspaces (which reduce to multiplication by a complex phase for eigenvalues of multiplicity one).

3. THE QUATERNIONIC EIGENVALUE PROBLEM

The quaternions $\mathbb{H}$ double the dimension of the complex numbers by adding two additional square roots of -1, usually denoted j and k. The multiplication table follows from

$$i^2 = j^2 = k^2 = -1 \qquad ij = k = -ji$$

and associativity; note that $\mathbb{H}$ is not commutative. Equivalently, $\mathbb{H}$ can be viewed via the Cayley-Dickson process as the sum of 2 copies of the complex numbers

$$\mathbb{H} = \mathbb{C} + \mathbb{C}j \tag{6}$$

with k being defined by $k = ij$.

The eigenvalue problem (1) for Hermitian matrices A over $\mathbb{H}$ immediately yields the first unexpected result: The eigenvalues need not be real. An example is given by

$$\begin{pmatrix} 1 & i \\ -i & 1 \end{pmatrix} \begin{pmatrix} 1 \\ k \end{pmatrix} = \begin{pmatrix} 1-j \\ k-i \end{pmatrix} = (1-j) \begin{pmatrix} 1 \\ k \end{pmatrix} \tag{7}$$

Furthermore, because of the lack of commutativity, multiples of eigenvectors are not necessarily eigenvectors. For instance, the vector

$$v_1 = \begin{pmatrix} \sqrt{2} \\ 1-i \end{pmatrix} \tag{8}$$

is an eigenvector of the matrix

$$A_1 = \begin{pmatrix} 0 & 1+i \\ 1-i & 0 \end{pmatrix} \tag{9}$$

with eigenvalue $\sqrt{2}$, but jv_1 is not an eigenvector of A_1. This example illustrates an important point: We must distinguish between right and left multiplication. Since

$$A(vq) = (Av)q \tag{10}$$

by associativity, *right* multiples of eigenvectors are indeed eigenvectors. For example, $v_1 j$ is an eigenvector of the matrix A_1 above, with the same eigenvalue ($\sqrt{2}$).

Similarly, we must carefully distinguish between the *left* eigenvalue problem (1) and the *right* eigenvalue problem

$$Av = v\lambda \tag{11}$$

It turns out that all the *right* eigenvalues of (quaternionic) Hermitian matrices are real.

Lemma $1_{\mathbb{H}}$: *The right eigenvalues of an $n \times n$ quaternionic Hermitian matrix $A = A^\dagger$ are real.*

Proof: This is just a careful rewrite of (2). Explicitly,

$$\overline{\lambda}(v^\dagger v) = (\overline{\lambda} v^\dagger)v = (Av)^\dagger v = (v^\dagger A)v = v^\dagger(Av) = v^\dagger(v\lambda) = (v^\dagger v)\lambda \tag{12}$$

and the result follows since $v^\dagger v \in \mathbb{R}$. ■

In the above, $\overline{\lambda}$ is now the quaternionic conjugate of λ and $\dagger$ denotes (quaternionic) Hermitian conjugation. Once the eigenvalues have been shown to be real, orthogonality of eigenvectors with different eigenvalues follows as in the complex case.

Lemma $2_{\mathbb{H}}$: *Right eigenvectors of an $n \times n$ quaternionic Hermitian matrix A corresponding to different eigenvalues are orthogonal.*

Proof: This is just a rewrite of (3). Explicitly,

$$\begin{aligned} \lambda_1(v_1^\dagger v_2) = (\lambda_1 v_1^\dagger)v_2 = (Av_1)^\dagger v_2 = (v_1^\dagger A)v_2 \\ = v_1^\dagger(Av_2) = v_1^\dagger(v_2\lambda_2) = (v_1^\dagger v_2)\lambda_2 \end{aligned} \tag{13}$$

so that, since $\lambda_m \in \mathbb{R}$, either $\lambda_1 = \lambda_2$ or $v_1^\dagger v_2 = 0$. ■

Finally, since the (right) eigenvalues are real, (right) multiples of eigenvectors are still eigenvectors.

Putting it all together, we obtain a decomposition of any (quaternionic) Hermitian matrix A of the form (4), where the real eigenvalues $\{\lambda_m\}$ and their eigenspaces are again unique.

Lemma $3_{\mathbb{H}}$: *For any $n \times n$ complex Hermitian matrix A, there exists an orthonormal basis of $\mathbb{H}^n$ consisting of eigenvectors of A.*

Theorem $1_{\mathbb{H}}$: *Let A be an $n \times n$ quaternionic Hermitian matrix. Then A can be expanded as in (4), where $\{v_m;\ m = 1, ..., n\}$ is an orthonormal basis of eigenvectors of A, with real eigenvalues λ_m.*

The proofs of each of these results is identical to the complex case previously considered. For eigenvalues of multiplicity one, the (unit) eigenvectors are unique up to a (right) quaternionic phase.

The right eigenvalue problem over $\mathbb{H}$ is therefore just a straightforward extension of the complex eigenvalue problem [3,4,5]. The left eigenvalue problem turns out to be of considerable interest as well, and will be considered elsewhere in the context of 2×2 octonionic



Figure 1: The representation of the octonionic multiplication table using the 7-point projective plane. Each of the 7 oriented lines represents a quaternionic triple.

Hermitian matrices [6] [3].

4. OCTONIONS AND OCTONIONIC MATRICES

a) The Octonions

The octonions $\mathbb{O}$ can be viewed via the Cayley-Dickson process as the direct sum of two copies of the quaternions [7]

$$\mathbb{O} = \mathbb{H} + \mathbb{H}\ell \tag{14}$$

where ℓ is yet another square root of -1. The octonions are thus spanned by the identity element 1 and the 7 imaginary units $\{i, j, k, \ell, i\ell, j\ell, k\ell\}$. These units can be grouped into associative "triples" in 7 different ways, each of which generates (the imaginary part of) a quaternionic subspace. Any three of these imaginary units which do not lie in a such a triple anti-associate. The multiplication table can be neatly summarized by appropriately labeling the 7-point projective plane, as shown in Figure 1. For a good introduction to the octonions, including some applications to physics, see [8,9].

When working with small numbers of octonions, it is important to realize that simplifications take place by virtue of the automorphism group G_2 of $\mathbb{O}$. For instance, a single

[3] Another argument that the right eigenvalue problem is the natural one is based on their use in diagonalizing a matrix. Cohn [5] considers quaternionic matrices A which are diagonalizable, in the sense that there exists an invertible matrix U and a diagonal matrix D such that

$$A = UDU^{-1}$$

But this is equivalent to

$$AU = UD$$

so that the columns of U are eigenvectors of A with right eigenvalues taken from the elements of D. We will return to this issue below. (Cohn uses the term *left eigenvalue* to describe the eigenvector problem for row vectors multiplied on the right by A; this is completely different from our use of the same term.)

octonion may be assumed without loss of generality to be complex, i.e. to lie in the complex subspace of $\mathbb{O}$ spanned by $\{1, i\}$. Similarly, a second octonion can be assumed to lie in the subspace spanned by $\{1, i, j\}$, and a third can be assumed to lie in the subspace spanned by $\{1, i, j, k, \ell\}$. Only when four or more octonions are involved is it necessary to consider "generic" octonions, involving all the basis directions.

The octonions are not associative. Nevertheless, since any 2 octonions lie in a quaternionic subspace, products involving only 2 different octonions (and their octonionic conjugates) do associate. For example,

$$p(pq) = p^2 q \tag{15}$$

which is a weak form of associativity known as *alternativity*.

The squared norm of an octonion a is given by

$$|a|^2 := a\overline{a} \tag{16}$$

where $\overline{a}$ denotes the (octonionic) conjugate of a. The commutator of a and b is given by

$$[a, b] := ab - ba \tag{17}$$

the associator of a, b, c is given by

$$[a, b, c] := (ab)c - a(bc) \tag{18}$$

and we use $A^\dagger$ to denote the (octonionic) Hermitian conjugate of the matrix A. Both the commutator and the associator are purely imaginary, totally antisymmetric, and change sign if any one of their arguments is replaced by its conjugate. Another octonionic product with the latter two properties is given by [4]

$$\Phi(a, b, c) = \frac{1}{2}\,\mathrm{Re}([a, \overline{b}]c) \tag{19}$$

b) Octonionic matrices

The lack of associativity complicates the treatment of matrices with octonionic entries. While matrix multiplication can be defined for matrices of arbitrary size in the usual way, only the 2×2 and 3×3 octonionic Hermitian matrices form Jordan algebras [12]. We therefore limit ourselves to these two cases.

Any complex number $x = x_1 + ix_2$ can be written as the real matrix $\begin{pmatrix} x_1 & -x_2 \\ x_2 & x_1 \end{pmatrix}$, which allows (complex) matrix multiplication to be rewritten in terms of real matrices. This process can be generalized to the quaternions, but fails for the octonions — as it must, since octonionic multiplication is not associative. But the complex number x can also be written as the real vector $\begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$, leading to a representation of (complex) multiplication as the product of a (real) matrix with a (real) vector, and this latter process does generalize to the octonions.

---

[4] $\Phi(a, b, c)$ is in fact the same as the *associative 3-form* [10,11]

$$\Phi(a, b, c) = \mathrm{Re}(a \times b \times c) = \frac{1}{2}\mathrm{Re}\Big(a(\overline{b}c) - c(\overline{b}a)\Big)$$

which reduces to the vector triple product when a, b, c are imaginary quaternions.
(Note that $a \times b \times c := \frac{1}{2}\big(a(\overline{b}c) - c(\overline{b}a)\big)$ is the triple cross product, *not* the iterated cross product.)

A similar process can be used to represent matrices over one of these division algebra as matrices over any smaller such algebra. Under this transformation a complex Hermitian matrix is mapped to a symmetric real matrix, and quaternionic and octonionic Hermitian matrices can be transformed into either real symmetric matrices or complex Hermitian matrices. For example, an $n \times n$ octonionic Hermitian matrix can be mapped to an $8n \times 8n$ symmetric real matrix (or a $4n \times 4n$ complex Hermitian matrix). It would seem as if we could therefore reduce the eigenvalue problem to the real case, but this is misleading for several reasons.

First of all, while the real formalism guarantees the existence of real eigenvalues, it does *not* rule out the possibility that eigenvalues might exist which are not real. This is because the general eigenvalue problem over a normed division algebra transforms into a *matrix* equation, which only reduces to an ordinary eigenvalue problem for real eigenvalues. As we will see, octonionic Hermitian matrices admit (right) eigenvalues which are not real.

Secondly, in the real formalism it is not very easy to determine the multiplicity of the real eigenvalues. One might expect n octonionic eigenvectors with at most n different real eigenvalues. The real formalism *does* guarantee us $8n$ independent (over $\mathbb{R}$) eigenvectors with real eigenvalues, but we are *not* guaranteed that the $8n$ eigenvalues occur with multiplicity 8 (or a multiple thereof). In fact, we will see below that this is not the case.

The final drawback of this approach is that the orthogonality of eigenvectors with different eigenvalues does *not* follow from the corresponding statement on the real, transformed eigenvectors. This is because the transformation does not preserve the inner product, which is real in one case and not in the other. Nevertheless, the real parts of the inner products do agree, so that

$$v^\dagger w + w^\dagger v = 0 \tag{20}$$

As we will see below, there is a stronger orthogonality condition on eigenvectors with different eigenvalues, which generalizes the usual notion of orthogonality.

For all of these reasons, we choose to work directly with the octonionic matrices.

5. 2×2 OCTONIONIC HERMITIAN MATRICES

Are the eigenvalues of octonionic Hermitian matrices real? Consider first the special case where A and v (but not necessarily λ) in (11) lie in a quaternionic subspace. Then (12) still holds, and alternativity allows us to shift the parentheses and conclude that λ is real.

For instance, since the diagonal entries of a 2×2 Hermitian matrix A are real, its components always lie in a *complex* subspace of $\mathbb{O}$, and therefore A clearly possesses eigenvectors which lie in the same complex subspace, and which have real eigenvalues. Multiplication (on the right) of these eigenvectors by an arbitrary octonion q leads to a new eigenvector, which lies in the quaternionic subspace spanned by the single octonionic direction in A and the octonionic multiple q. Furthermore, this new eigenvector has the same real eigenvalue as the original eigenvector, that is

$$Av = v\lambda \Longrightarrow A(vq) = (Av)q = (v\lambda)q = (vq)\lambda \tag{21}$$

since A and v are complex and λ is real.

In general, however, the key use of associativity in the middle of the derivation of (12) is not allowed. We are thus led to suspect that there exist octonionic Hermitian matrices

which admit (right) eigenvalues which are *not* real. This turns out to be correct, as is shown by the following example:

$$\begin{pmatrix} 1 & i \\ -i & 1 \end{pmatrix} \begin{pmatrix} j \\ \ell \end{pmatrix} = \begin{pmatrix} j+i\ell \\ \ell - k \end{pmatrix} = \begin{pmatrix} j \\ \ell \end{pmatrix} (1 - k\ell) \tag{22}$$

Further details for the case of octonionic Hermitian matrices whose eigenvalues are not real will appear elsewhere [6].

a) The real eigenvalue problem

We now turn to the case of *real* eigenvalues. The general 2×2 octonionic Hermitian matrix can be written

$$A = \begin{pmatrix} p & a \\ \overline{a} & m \end{pmatrix} \tag{23}$$

with $p, m \in \mathbb{R}$ and $a \in \mathbb{O}$, and satisfies its characteristic equation

$$A^2 - (\operatorname{tr} A)\, A + (\det A)\, I = 0 \tag{24}$$

where $\operatorname{tr} A$ denotes the trace of A, and where there is no difficulty defining the determinant of A as usual via

$$\det A = pm - |a|^2 \tag{25}$$

If $a = 0$ the eigenvalue problem is trivial, so we assume $a \neq 0$. If we set

$$v = \begin{pmatrix} x \\ y \end{pmatrix} \tag{26}$$

then (11) leads to

$$y = \frac{\overline{a}x}{|a|^2}\,(\lambda - p) \qquad\qquad x = \frac{ay}{|a|^2}\,(\lambda - m) \tag{27}$$

from which it follows (unless $v = 0$) that

$$\det(\lambda I - A) = \lambda^2 - (\operatorname{tr} A)\,\lambda + \det A = 0 \tag{28}$$

as usual. Eigenvectors can thus be written in either of the forms

$$v = \begin{pmatrix} |a|^2 \\ \overline{a}(\lambda - p) \end{pmatrix} \xi \qquad\qquad v = \begin{pmatrix} a(\lambda - m) \\ |a|^2 \end{pmatrix} \xi \tag{29}$$

where $\xi \in \mathbb{O}$ is arbitrary and where λ is either of the 2 solutions of (28). [5] This shows that *all* eigenvectors of 2×2 Hermitian matrices with real eigenvalues are obtained from the usual complex eigenvectors by (right) multiplication by an arbitrary octonion.

Lemma $\mathbf{1}_{\mathbb{O}_2}$: *Let A be a 2×2 complex Hermitian matrix. Then w is an octonionic eigenvector of A with real eigenvalue λ if and only if $w = v\xi$, where $\xi \in \mathbb{O}$ is arbitrary and where v is a complex eigenvector of A with the same eigenvalue.*

Since a 2×2 octonionic Hermitian matrix contains only one independent octonion, we can assume without loss generality that any such matrix is complex, and thus apply this lemma to it.

[5] Note that $x \neq 0 \neq y$ if $a \neq 0$, and that in this case the 2 solutions of (28) are distinct.

b) Orthogonality and Decompositions

As already noted, since A lies in a complex subspace of $\mathbb{O}$, it admits a complete set of complex eigenvectors with real eigenvalues, which can be used to obtain the decomposition (4). But what about a decomposition in terms of the general solution of the real eigenvalue problem?

The general solution is given by (29). Choosing the first form, we obtain a complete set of eigenvectors by considering both solutions $\lambda_\pm$ to (28), obtaining

$$v_\pm = \begin{pmatrix} |a|^2 \\ \overline{a}(\lambda_\pm - p) \end{pmatrix} \xi_\pm \tag{30}$$

Are these eigenvectors orthogonal? Direct computation yields

$$v_+^\dagger v_- = -|a|^2 a \left[a, \xi_+, \xi_-\right] \tag{31}$$

where we have used

$$a[a,x,y] = -a[\overline{a},y,\overline{x}] \equiv [a,\overline{a}y,\overline{x}] = [\overline{x},a,\overline{a}y] = (\overline{x}a)(\overline{a}y) - |a|^2\overline{x}y \tag{32}$$

or equivalently

$$a\Big((\overline{ax})y\Big) = (\overline{x}a)(\overline{a}y) \tag{33}$$

for any octonions a, x, y. Thus, the eigenvectors are not necessarily orthogonal in the traditional sense except for the quaternionic eigenvalue problem, when the associator automatically vanishes.

At first sight, this apparent lack of orthogonality seems to rule out a decomposition of the form (4). However, due to the lack of associativity, what is needed for (4) to hold is an appropriately generalized notion of orthogonality, namely

Definition: *Let v and w be two octonionic vectors. We will say that w is orthogonal to v if*

$$(vv^\dagger)\, w = 0 \tag{34}$$

The vectors $\{v, w\}$ are orthonormal if in addition $v^\dagger v = 1 = w^\dagger w$.

Direct computation shows that the eigenvectors $v_\pm$ above are indeed mutually orthogonal in this sense, which provides a computational proof of the following lemma.

Lemma $\mathbf{2}_{\mathbb{O}_2}$: *If v and w are eigenvectors of the 2×2 octonionic Hermitian matrix A corresponding to different real eigenvalues, then v and w are mutually orthogonal in the sense of* (34).

Proof: From Lemma $1_{\mathbb{O}_2}$, we can write

$$v = \hat{v}\alpha, \quad w = \hat{v}\beta \qquad (\hat{v}, \hat{w} \in \mathbb{C}; \alpha, \beta \in \mathbb{O})$$

where $\mathbb{C} \subset \mathbb{O}$ is the complex subspace containing the elements of A. But then $vv^\dagger = |\alpha|^2 \hat{v}\hat{v}^\dagger$, and

$$(vv^\dagger)w = |\alpha|^2(\hat{v}\hat{v}^\dagger)(\hat{w}\beta)$$

which associates since only 2 octonionic directions are involved. But $\hat{v}^\dagger\hat{w} = 0$ by the usual properties of complex eigenvectors. ■

In order for a decomposition of the form (4) to exist, we also need a vector version of alternativity, which in fact holds for octonionic vectors of any size: [6]

Proposition 1: *For any octonionic vector* $v \in \mathbb{O}^n$,

$$(vv^\dagger)\, v = v\, (v^\dagger v) \tag{35}$$

This proposition shows in particular that any normalized vector v is an eigenvector of the matrix $vv^\dagger$ with eigenvalue 1, as required by (4). We conclude that the decomposition (4) holds unchanged for real eigenvalues. We thus have:

Theorem $1_{\mathbb{O}_2}$: *Let A be a 2×2 octonionic Hermitian matrix. Then A can be expanded as in (4), where $\{v_1, v_2\}$ are orthonormal (as per (34)) eigenvectors of A corresponding to the real eigenvalues λ_m.*

Proof: Provided the real eigenvalues of A are distinct, Lemma $2_{\mathbb{O}_2}$ guarantees the existence of orthonormal eigenvectors, which are also eigenvectors of the decomposition (4) with the same eigenvalues, and the result follows. But if A has a repeated eigenvalue, it must be a multiple of the identity matrix, for which the result holds. ■

Using the same technique as in Lemma $2_{\mathbb{O}_2}$, it is straightforward to show that

$$(vv^\dagger)(vv^\dagger) = (v^\dagger v)(vv^\dagger) \tag{36}$$

for any $v \in \mathbb{O}^2$, and that

$$(vv^\dagger)(ww^\dagger) = 0 \tag{37}$$

if v and w are eigenvectors of A with distinct real eigenvalues, since each term in parentheses lies in $\mathbb{C}$. The decomposition in the preceding theorem is thus in terms of orthonormal idempotents $v_i v_i^\dagger$. We also have

$$\frac{vv^\dagger}{v^\dagger v} + \frac{ww^\dagger}{w^\dagger w} = I \tag{38}$$

which could be proved directly using the fact that the left-hand side has repeated eigenvalue 1. Furthermore, since by Lemma $2_{\mathbb{O}_2}$ A and v contain only 2 octonionic directions,

$$(Av)v^\dagger = A(vv^\dagger) \tag{39}$$

which leads to the following one-line alternative derivation of Theorem $1_{\mathbb{O}_2}$

$$A = A\left(\sum_{i=1}^{2} v_i v_i^\dagger\right) = \sum_{i=1}^{2} A(v_i v_i^\dagger) = \sum_{i=1}^{2} (Av_i) v_i^\dagger = \sum_{i=1}^{2} \lambda_i v_i v_i^\dagger \tag{40}$$

As we will see below, however, this argument relies heavily on Lemma $2_{\mathbb{O}_2}$, which fails in the 3×3 case.

We have not yet discussed whether the orthonormal eigenvectors in the preceding theorem constitute a basis of $\mathbb{O}^2$. For any orthonormal vectors, we have the following lemma.

[6] For 2-component vectors, this proposition is just the 3-Ψ's rule of supersymmetry theory [13].

Lemma $3_{\mathbb{O}_2}$: *Let $v, w \in \mathbb{O}^2$ be orthonormal in the sense of (34), and let g be any vector in $\mathbb{O}^2$. Then*

$$g = (vv^\dagger)\, g + (ww^\dagger)\, g$$

Proof: This follows immediately from (38). ■

In the associative case, this lemma shows how to write any vector g in terms of its components along v and w, thus establishing $\{v, w\}$ as a basis. One could adopt similar language in the nonassociative case, although the "component" of g "along" v would no longer point in the v direction. Nevertheless, this terminology is extremely attractive, as it allows the Gram-Schmidt orthogonalization process to be used to determine the component of one vector orthogonal to another.

Proposition 2: *Let $v, w \in \mathbb{O}^2$. Then*

$$(vv^\dagger)\left(w - \frac{(vv^\dagger)}{v^\dagger v}\, w\right) = 0 \tag{41}$$

Proof: This follows from the alternativity of 2×2 octonionic Hermitian matrices and (36). ■

6. 3×3 OCTONIONIC HERMITIAN MATRICES

We now turn to the 3×3 case. It is not immediately obvious that 3×3 octonionic Hermitian matrices have a well-defined determinant, let alone a characteristic equation. We therefore first review some of the properties of these matrices before turning to the eigenvalue problem. As in the 2×2 case, over the octonions there will be solutions of the eigenvalue problem with eigenvalues which are not real; we consider here only the real eigenvalue problem.

a) Jordan matrices

The 3×3 octonionic Hermitian matrices, henceforth referred to as *Jordan matrices*, form the exceptional Jordan algebra (also called the Albert algebra) under the Jordan product [7]

$$A \circ B := \frac{1}{2}(AB + BA) \tag{42}$$

which is commutative, but not associative. A special case of this is

$$A^2 \equiv A \circ A \tag{43}$$

and we *define*

$$A^3 := A^2 \circ A = A \circ A^2 \tag{44}$$

Remarkably, with these definitions, Jordan matrices satisfy the usual characteristic equation [10]

$$A^3 - (\mathrm{tr}\, A)\, A^2 + \sigma(A)\, A - (\det A)\, I = 0 \tag{45}$$

where $\sigma(A)$ is defined by

$$\sigma(A) := \frac{1}{2}\left((\mathrm{tr}\, A)^2 - \mathrm{tr}\,(A^2)\right) \tag{46}$$

[7] The 2×2 octonionic Hermitian matrices form a special Jordan algebra since they are alternative [12].

and where the determinant of A is defined abstractly in terms of the Freudenthal product. [8] Concretely, if

$$A = \begin{pmatrix} p & a & \overline{b} \\ \overline{a} & m & c \\ b & \overline{c} & n \end{pmatrix} \tag{47}$$

with $p, m, n \in \mathbb{R}$ and $a, b, c \in \mathbb{O}$ then

$$\begin{aligned} \operatorname{tr} A &= p + m + n \\ \sigma(A) &= pm + pn + mn - |a|^2 - |b|^2 - |c|^2 \\ \det A &= pmn + b(ac) + \overline{b(ac)} - n|a|^2 - m|b|^2 - p|c|^2 \end{aligned} \tag{48}$$

b) The real eigenvalue problem

As discussed above, $n \times n$ Hermitian matrices over any of the normed division algebras can be rewritten as symmetric $kn \times kn$ real matrices, where k denotes the dimension of the underlying division algebra, it is clear that a 3×3 octonionic Hermitian matrix must have $8 \times 3 = 24$ real eigenvalues [1]. However, as we now show, instead of having (a maximum of) 3 distinct real eigenvalues, each with multiplicity 8, we show that there are (a maximum of) 6 distinct real eigenvalues, each with multiplicity 4.

The reason for this is that, somewhat surprisingly, a (real) eigenvalue λ of a Jordan matrix A does *not* in general satisfy the characteristic equation (45). [9] To see this, consider the eigenvalue equation (1), with A as in (47), $\lambda \in \mathbb{R}$, and where

$$v = \begin{pmatrix} x \\ y \\ z \end{pmatrix} \tag{49}$$

Explicit computation yields

$$(\lambda - p)x = ay + \overline{b}z \tag{50}$$

$$(\lambda - m)y = cz + \overline{a}x \tag{51}$$

$$(\lambda - n)z = bx + \overline{c}y \tag{52}$$

so that

$$(\lambda - p)(\lambda - m)y = (\lambda - p)(cz + \overline{a}x) = (\lambda - p)cz + \overline{a}(ay + \overline{b}z) \tag{53}$$

---

[8] The Freudenthal product of two Jordan matrices A and B is given by [14]

$$A * B = A \circ B - \frac{1}{2}\Big(A \operatorname{tr}(B) + B \operatorname{tr}(A)\Big) + \frac{1}{2}\Big(\operatorname{tr}(A) \operatorname{tr}(B) - \operatorname{tr}(A \circ B)\Big)$$

The determinant can then be defined as

$$\det(A) = \frac{1}{3} \operatorname{tr}\Big((A * A) \circ A\Big)$$

[9] Ogievetskiĭ [2] constructed a 6th order polynomial satisfied by the real eigenvalues, which he called the characteristic equation. This polynomial is presumably equivalent to the modified characteristic equations (for both values of r) given below.

which implies

$$\Big[(\lambda-p)(\lambda-m)-|a|^2\Big]\,y=\overline{a}(\overline{b}z)+(\lambda-p)cz \tag{54}$$

Assume first that $\lambda\neq p$. Using (50) and (54) in (52) leads to

$$\begin{aligned}\Big[(\lambda-p)(\lambda-m)-|a|^2\Big](\lambda-p)(\lambda-n)z&\\ =\Big[(\lambda-p)(\lambda-m)-|a|^2\Big]\,(\lambda-p)(bx+\overline{c}y)&\\ =\Big[(\lambda-p)(\lambda-m)-|a|^2\Big]\,b(ay+\overline{b}z)+(\lambda-p)\overline{c}\Big[\overline{a}(\overline{b}z)+(\lambda-p)cz\Big]&\\ =b\Big[a\Big(\overline{a}(\overline{b}z)+(\lambda-p)cz\Big)\Big]+\Big[(\lambda-p)(\lambda-m)-|a|^2\Big]\,b(\overline{b}z)&\\ \qquad+(\lambda-p)\overline{c}\Big[\overline{a}(\overline{b}z)+(\lambda-p)cz\Big]&\\ =(\lambda-p)\Big[(\lambda-m)|b|^2z+(\lambda-p)|c|^2z+b\Big(a(cz)\Big)+\overline{c}\,\Big(\overline{a}(\overline{b}z)\Big)\Big]&\end{aligned} \tag{55}$$

Expanding this out and comparing with (48) results finally in [10]

$$\begin{aligned}\Big[\det(\lambda I-A)\Big]z\equiv\Big[\lambda^3-(\operatorname{tr}A)\,\lambda^2+\sigma(A)\,\lambda-\det A\Big]\,z&\\ =b\Big(a(cz)\Big)+\overline{c}\,\Big(\overline{a}(\overline{b}z)\Big)-\Big[b(ac)+(\overline{c}\,\overline{a})\overline{b}\Big]\,z&\end{aligned} \tag{56}$$

Now consider the case $\lambda=p$. We still have (54), which here takes the form

$$-|a|^2y=\overline{a}(\overline{b}z) \tag{57}$$

Inserting this into (51), we can solve for x, obtaining

$$-|a|^2x=a(cz)+(p-m)\overline{b}z \tag{58}$$

Finally, inserting (57) and (58) in (52) yields

$$-\Big(|a|^2(p-n)+|b|^2(p-m)\Big)\,z=b\Big(a(cz)\Big)+\overline{c}\,\Big(\overline{a}(\overline{b}z)\Big) \tag{59}$$

Comparing with (48) and using $\lambda=p$, we see that (56) still holds, and thus holds in general.

If a, b, c, and z associate, the RHS of (56) vanishes, and λ does indeed satisfy the characteristic equation (45); this will not happen in general. However, since the LHS of (56) is a real multiple of z, this must also be true of the RHS, so that

$$b\Big(a(cz)\Big)+\overline{c}\,\Big(\overline{a}(\overline{b}z)\Big)-\Big[b(ac)+(\overline{c}\,\overline{a})\overline{b}\Big]\,z=rz\qquad\qquad r\in\mathbb{R} \tag{60}$$

which can be solved to yield a quadratic equation for r as well as constraints on z.

Lemma $\mathbf{1}_{\mathbb{O}_3}$: *The real eigenvalues of the 3×3 octonionic Hermitian matrix A satisfy the modified characteristic equation*

$$\det(\lambda I-A)=\lambda^3-(\operatorname{tr}A)\,\lambda^2+\sigma(A)\,\lambda-\det A=r \tag{61}$$

where r is either of the two roots of

$$r^2+4\Phi(a,b,c)\,r-\Big|[a,b,c]\Big|^2=0 \tag{62}$$

with a,b,c as defined by (47) *and where Φ was defined in* (19).

[10] We have recently been able to generalize this to the case where λ is not real [6].

Proof: These results were obtained using *Mathematica* to solve (60) by brute force for real r and octonionic z given generic octonions a, b, c [15]. ■

Furthermore, provided that $[a,b,c] \neq 0$, each of x, y, and z can be shown to admit an expansion in terms of 4 real parameters.

Corollary 1: *With A and r as above, and assuming $[a,b,c] \neq 0$,*

$$z = (\alpha a + \beta b + \gamma c + \delta)\left(1 + \frac{[a,b,c]\,r}{\Big|[a,b,c]\Big|^2}\right) \tag{63}$$

with $\alpha, \beta, \gamma, \delta \in \mathbb{R}$. Similar expansions hold for x and y.

The real paramaters $\alpha, \beta, \gamma, \delta$ may be freely specified for one (nonzero) component, say z; the remaining components x, y have a similar form which is then fully determined by (50)–(52).

Corollary 2: *The real eigenvalues of $\overline{A}$ are the same as those of A.*

Proof: Direct computation (or (48)) shows that

$$\det \overline{A} = \det A - 4\Phi(a,b,c) \tag{64}$$

But $-4\Phi(a,b,c)$ is precisely the sum of the roots of (62), and replacing A by $\overline{A}$ merely flips the sign of r, that is $r[\overline{A}] = -r[A]$. Thus, the 2 possible values of $\det A + r[A]$ are precisely the same as those for $\det \overline{A} + r[\overline{A}]$. Since $\operatorname{tr} \overline{A} = \operatorname{tr} A$ and $\sigma(\overline{A}) = \sigma(A)$, (61) is unchanged. ■

The solutions of (61) are real, since the corresponding 24×24 real symmetric matrix has 24 real eigenvalues. We will refer to the 3 real solutions of (61) corresponding to a single value of r as a *family* of eigenvalues of A. There are thus 2 families of real eigenvalues, each corresponding to 4 independent (over $\mathbb{R}$) eigenvectors.

We note several intriguing properties of these results. If A is in fact complex, then the only solution of (62) is $r = 0$, and we recover the usual characteristic equation with a unique set of 3 (real) eigenvalues. If A is quaternionic, then one solution of (62) is $r = 0$, leading to the standard set of 3 real eigenvalues and their corresponding quaternionic eigenvectors. However, unless a, b, c involve only two independent imaginary quaternionic directions (in which case $\Phi(a,b,c) = 0 = [a,b,c]$), there will also be a nonzero solution for r, leading to a second set of 3 real eigenvalues. From the preceding corollary, we see that this second set of eigenvalues consists precisely of the usual ($r = 0$) eigenvalues of $\overline{A}$! Furthermore, since $A(\ell v) = \ell(\overline{A}v)$ for $A, v \in \mathbb{H}$ and imaginary $\ell \in \mathbb{O}$ orthogonal to $\mathbb{H}$, the eigenvectors of A corresponding to $r \neq 0$ are precisely ℓ times the quaternionic ($r = 0$) eigenvectors of $\overline{A}$. In this sense, the octonionic eigenvalue problem for quaternionic A is equivalent to the quaternionic eigenvalue problem for both A and $\overline{A}$ together. Finally, if A is octonionic (so that in particular $[a,b,c] \neq 0$), then there are two distinct solutions for r, and hence two different sets of real eigenvalues, with corresponding eigenvectors. Note that if $\det A = 0 \neq [a,b,c]$ then all of the eigenvalues of A will be nonzero!

The final suprise lies with the orthogonality condition for eigenvectors v, w corresponding to different eigenvalues. It is *not* true (cf. (31)) that $v^\dagger w = 0$, although the real part (20) of

this expression does vanish. However, just as in the 2×2 case, what is needed to ensure a decomposition of the form (4) is (34), and a lengthy, direct computation verifies that (34) holds *provided that* both eigenvectors correspond to the same value of r.

Lemma $\mathbf{2}_{\mathbb{O}_3}$: *If v and w are eigenvectors of the 3×3 octonionic Hermitian matrix A corresponding to different real eigenvalues in the same family (same r value), then v and w are mutually orthogonal in the sense of* (34)*.*

Proof: The modified characteristic equation (61) can be used to eliminate cubic and higher powers of λ from any expression. Furthermore, given two distinct eigenvalues $\lambda_1 \neq \lambda_2$, subtracting the two versions of (61) and factoring the result leads to the equation

$$(\lambda_1^2 + \lambda_1\lambda_2 + \lambda_2^2) - \mathrm{tr}\, A(\lambda_1 + \lambda_2) + \sigma(A) = 0 \tag{65}$$

which can be used to eliminate quadratic terms in one of the eigenvalues. [11] ■

For Jordan matrices, we thus obtain *two* decompositions of the form (4), corresponding to the two sets of real eigenvalues. For each, the eigenvectors are fixed up to orthogonal transformations which preserve the form (63) of z.

Theorem $\mathbf{1}_{\mathbb{O}_3}$: *Let A be a 3×3 octonionic Hermitian matrix. Then A can be expanded as in* (4)*, where $\{v_1, v_2, v_3\}$ are orthonormal (as per* (34)*) eigenvectors of A corresponding to the real eigenvalues λ_m, which belong to the same family (same r value).*

Proof: Fix a family of real eigenvalues of A by fixing r. If the eigenvalues are distinct, then the previous theorem guarantees the existence of orthonormal eigenvectors, which are also eigenvectors of the decomposition (4) with the same eigenvalues, and the result follows.

If the eigenvalues are the same, the family consists of a single real eigenvalue λ with multiplicity 3. Then $\mathrm{tr}\,(A) = 3\lambda$ and $\sigma(A) = 3\lambda^2$. Writing out these two equations in terms of the components (47) of A, and inserting the first into the second, results in a quadratic equation for λ; the discriminant D of this equation satisfies $D \leq 0$. But λ is assumed to be real, which forces $D = 0$, which in turn forces A to be a multiple of the identity matrix, for which the result holds.

The remaining case is when one eigenvalue, say μ, has multiplicity 2 and one has multiplicity 1. Letting v be a (normalized) eigenvector with eigenvalue μ, consider the matrix

$$X = A - \alpha\, vv^\dagger \tag{66}$$

[11] We used *Mathematica* to implement these simplifications in a brute force verification of (34) in this context, which ran for 6 hours on a SUN Sparc20 with 224 Mb of RAM [15].

with $\alpha \in \mathbb{R}$. For most values of α, X will have 3 distinct real eigenvalues, whose eigenvectors will be orthogonal by the previous theorem. But this means that eigenvectors of X are also eigenvectors of A; the required decomposition of A is obtained from that of X simply by solving (66) for A. ■

Note in particular that for some quaternionic matrices with determinant equal to zero, one and only one of these two decompositions will contain the eigenvalue zero.

In the 2×2 case, (36) tells us that, for normalized v, $vv^\dagger$ squares to itself, and hence is idempotent. As already noted, the decomposition of Theorem $1_{\mathbb{O}_2}$ is thus an idempotent decomposisition. But (36) fails in the 3×3 case, so that the decomposition in Theorem $1_{\mathbb{O}_3}$ is therefore *not* an idempotent decomposition.

It is nevertheless straightforward to show, in analogy with (38), that if u, v, and w are orthonormal in the sense of (34), then

$$uu^\dagger + vv^\dagger + ww^\dagger = I \tag{67}$$

since the left-hand side has eigenvalue 1 with multiplicity 3. This permits us to view $\{u, v, w\}$ as a basis of $\mathbb{O}^3$ in the following sense

Lemma $3_{\mathbb{O}_3}$: *Let $u, v, w \in \mathbb{O}^3$ be orthonormal in the sense of (34), and let g be any vector in $\mathbb{O}^3$. Then*

$$g = (uu^\dagger)\, g + (vv^\dagger)\, g + (ww^\dagger)\, g \tag{68}$$

Proof: This follows immediately from (67). ■

However, another consequence of the failure of (36) in the 3×3 case is that the Gram-Schmidt orthogonalization procedure (41) no longer works. It appears to be fortuitous that we are nevertheless able to find orthonormal eigenvectors in the 3×3 case with repeated eigenvalues; we suspect that this might fail in general, perhaps already in the 4×4 case with an eigenvalue of multiplicity 3.

7. DISCUSSION

Our main result is (61) together with (62), which shows how to modify the characteristic equation for a 3×3 octonionic Hermitian matrix in order to find its real eigenvalues, and which further shows that there are in general 2 families of solutions of these equations, each consisting of 3 eigenvalues of multiplicity 4.

We have further shown how to use the corresponding eigenvectors to recover the usual decomposition (4) of a Hermitian matrix in terms of its eigenvectors for both 2×2 and 3×3 octonionic Hermitian matrices. In the process we were led to introduce an appropriately generalized notion of orthogonality, namely (34).

We can relate our notion of orthonormality to the usual one by noting that n vectors in $\mathbb{O}^n$ which are orthonormal in the sense (34) satisfy

$$vv^\dagger + ... + ww^\dagger = I \tag{69}$$

If we define a matrix U whose columns are just $v, ..., w$, then this statement is equivalent to

$$UU^\dagger = I \tag{70}$$

Over the quaternions, left matrix inverses are the same as right matrix inverses, and we would also have

$$U^\dagger U = I \tag{71}$$

or equivalently

$$v^\dagger v = 1 = ... = w^\dagger w; \qquad v^\dagger w = 0 = ... \tag{72}$$

which is just the standard notion of orthogonality. These two notions of orthogonality fail to be equivalent over the octonions; we have been led to view the former as more fundamental.

We can now rewrite the eigenvalue equation (11) in the form

$$AU = UD \tag{73}$$

where D is a diagonal matrix whose entries are the real eigenvalues. Multiplying (73) on the left by $U^\dagger$ yields

$$U^\dagger(AU) = U^\dagger(UD) = (U^\dagger U)D \tag{74}$$

(since D is real), but this does not lead to a diagonalization of A since, as noted above, $U^\dagger U$ is not in general equal to the identity matrix. However, Theorem $1_{\mathbb{O}_3}$ can be rewritten as

$$A = UDU^\dagger \tag{75}$$

so that in this sense A is diagonalizable. Furthermore, multiplication of (73) on the right by $U^\dagger$ shows that

$$(AU)U^\dagger = (UD)U^\dagger = A = A(UU^\dagger) \tag{76}$$

and this assertion of associativity can be taken as a restatement of both Theorem $1_{\mathbb{O}_2}$ and Theorem $1_{\mathbb{O}_3}$. In the 2×2 case, this associativity holds for a single eigenvector v, which is (39), and which was used in the one-line proof (40) of Theorem $1_{\mathbb{O}_2}$. However, (39) fails in the 3×3 case, and we are unaware of a correspondingly elegant proof of Theorem $1_{\mathbb{O}_3}$.

Many of our results were obtained using *Mathematica* [15]. In fact, the *only* proof we currently have of the 3×3 orthogonality result, namely Lemma $2_{\mathbb{O}_3}$, uses *Mathematica* to explicitly perform a horrendous, but exact, algebraic computation. While one could hope for a more elegant mathematical proof of this result, the *Mathematica* computation nevertheless establishes a result which would otherwise remain for the moment merely a conjecture. This is a good example of being able to use the computer to verify one's intuition when it may not be possible to do so otherwise. This issue is further discussed in [15].

Finally, it is intriguing that (some) Hermitian octonionic matrices admit eigenvalues which are not real. In particular, this means that octonionic self-adjoint operators do not necessarily have a (purely) real spectrum. We plan to report separately on these matters [6].

ACKNOWLEDGMENTS

We are grateful to Al Agnew for unwittingly suggesting the topic, and to Tom Craven, Burt Fein, and Tony Sudbery for discussions.

REFERENCES

1. H. H. Goldstine & L. P. Horwitz, *On a Hilbert Space with Nonassociative Scalars*, Proc. Nat. Aca. **48**, 1134 (1962).
2. O. V. Ogievetskiĭ, *A Characteristic Equation for* 3×3 *Matrices over the Octonions*, Uspekhi Mat. Nauk **36**, 197–198 (1981); reviewed in Mathematical Reviews 83e:15017.
3. H. C. Lee, *Eigenvalues and Canonical Forms of Matrices with Quaternion Coefficients*, Proc. Roy. Irish Acad. **A52**, 253–260 (1949).
4. J. L. Brenner, *Matrices of Quaternions*, Pacific J. Math. **1**, 329–335 (1951).
5. P. M. Cohn, **Skew Field Constructions**, Cambridge University Press, 1977.
6. Tevian Dray, Jason Janesky, and Corinne A. Manogue, *Octonionic Hermitian Matrices with Non-Real Eigenvalues* (in preparation).
7. Richard D. Schafer, **An Introduction to Nonassociative Algebras**, Academic Press, New York, 1966 & Dover, Mineola NY, 1995.
8. S. Okubo, **Introduction to Octonion and Other Non-Associative Algebras in Physics**, Cambridge University Press, Cambridge, 1995.
9. Feza Gürsey and Chia-Hsiung Tze, **On the Role of Division, Jordan and Related Algebras in Particle Physics**, World Scientific, Singapore, 1996.
10. F. Reese Harvey, **Spinors and Calibrations**, Academic Press, Boston, 1990.
11. G. B. Gureirch, **Foundations of the Theory of Algebraic Invariants**, P. Noordhoff, Groningen, 1964.
12. Nathan Jacobson, **Structure and Representations of Jordan Algebras**, Amer. Math. Soc. Colloq. Publ. **39**, American Mathematical Society, Providence, 1968.
13. Corinne A. Manogue & Anthony Sudbery, *General Solutions of Covariant Superstring Equations of Motion*, Phys. Rev. **D40**, 4073-4077 (1989);
    Tevian Dray and Corinne A. Manogue, *Associators and the 3-Ψ's Rule*, (in preparation).
14. P. Jordan, J. von Neumann, and E. Wigner, Ann. Math. **36**, 29 (1934);
    H. Freudenthal, Adv. Math. **1**, 145 (1964).
15. Tevian Dray and Corinne A. Manogue, *Finding Octonionic Eigenvectors Using Mathematica*, Comput. Phys. Comm. (invited paper; submitted).